

The *Caenorhabditis elegans* Centrosomal Protein SPD-2 Is Required for both Pericentriolar Material Recruitment and Centriole Duplication

Laurence Pelletier,^{1,*} Nurhan Özlü,¹ Eva Hannak,^{1,2} Carrie Cowan,¹ Bianca Habermann,³ Martine Ruer,¹ Thomas Müller-Reichert,¹ and Anthony A. Hyman¹

¹Max Planck Institute of Molecular Cell Biology
and Genetics
Pfortenhauerstrasse 108
01307 Dresden
Germany

*Correspondence: pelletie@mpi-cbg.de

²Present address: Department of Molecular and Cell Biology, University of California, Berkeley, 315 LSA, Berkeley, CA 94720-3200.

³Present address: Scionics Computer Innovation, Pfortenhauerstrasse 110, 01307 Dresden, Germany.

(Current Biology 14, 863–873; May 25, 2004)

The sequences of the oligos used to generate SPD-2 dsRNA as indicated in the online supplementary material accompanying this paper were incorrect. The correct sequences of the primers used are as follows (T3 and T7 promoter sequences are underlined):

Forward:

aattaccctcactaaaggTGCATGCGAATAAGACGAAG

Reverse:

taatcgactcactataggTTGCGGACACAGAAAACAAA

This error does not change any of the results presented in the paper.

The authors would like to apologize to their colleagues for any inconvenience this may have caused.

DOI: 10.1016/j.cub.2006.06.021
